



Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective

Matthew Baumgart^a, Heather M. Snyder^{b,*}, Maria C. Carrillo^b, Sam Fazio^c,
Hye Kim^a, Harry Johns^d

^aDivision of Public Policy, Alzheimer's Association, Washington, DC, USA

^bDivision of Medical & Scientific Relations, Alzheimer's Association, Chicago, IL, USA

^cDivision of Constituent Relations, Alzheimer's Association, Chicago, IL, USA

^dPresident & CEO, Alzheimer's Association, Chicago, IL, USA

Abstract

An estimated 47 million people worldwide are living with dementia in 2015, and this number is projected to triple by 2050. In the absence of a disease-modifying treatment or cure, reducing the risk of developing dementia takes on added importance. In 2014, the World Dementia Council (WDC) requested the Alzheimer's Association evaluate and report on the state of the evidence on modifiable risk factors for cognitive decline and dementia. This report is a summary of the Association's evaluation, which was presented at the October 2014 WDC meeting. The Association believes there is sufficient evidence to support the link between several modifiable risk factors and a reduced risk for cognitive decline, and sufficient evidence to suggest that some modifiable risk factors may be associated with reduced risk of dementia. Specifically, the Association believes there is sufficiently strong evidence, from a population-based perspective, to conclude that regular physical activity and management of cardiovascular risk factors (diabetes, obesity, smoking, and hypertension) reduce the risk of cognitive decline and may reduce the risk of dementia. The Association also believes there is sufficiently strong evidence to conclude that a healthy diet and lifelong learning/cognitive training may also reduce the risk of cognitive decline.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords:

World Dementia Council; Alzheimer's Association; Alzheimer's disease; Cognitive decline; Dementia; Risk factors; Modifiable risk factors; Cardiovascular disease risk factors; Lifestyle risk factors; Physical activity; Diabetes; Obesity; Smoking; Hypertension; Diet; Lifelong learning; Cognitive training

1. Introduction

An estimated 47 million people worldwide are living with dementia in 2015 [1], and this number is projected to triple by 2050 [2]. In the absence of a disease-modifying treatment or cure, reducing the risk of developing dementia takes on added importance. Even when effective treatments become available, risk reduction will likely remain a fundamental strategy in reducing the number of individuals affected; for many non-communicable diseases with available treatments

(such as diabetes, cancer, and heart disease), risk reduction efforts remain a major component of the campaigns against these diseases.

As a science-based advocacy organization, the Alzheimer's Association—the largest voluntary health organization dedicated to Alzheimer's disease and other dementias—is the global nonprofit leader in Alzheimer's disease research and the leading resource for more than 5 million individuals living with the disease in the United States and their caregivers. In this role, we are often asked for both expertise and guidance related to risk reduction for Alzheimer's disease. The Association monitors the science and develops its positions accordingly.

*Corresponding author. Tel.: 312-335-5184; Fax: 866-875-2553.

E-mail address: hsnyder@alz.org

In December 2013, the G8 nations—Canada, France, Germany, Italy, Japan, Russia, the United Kingdom, and the United States—created the World Dementia Council (WDC) [3] to provide global advocacy and leadership on key dementia challenges. The WDC is composed of individuals from around the world with a wide range of expertise and from a wide range of disciplines. One of the WDC's priority areas is potential risk reduction, both in the absence of treatments and after the time at which a treatment or treatments become available. However, the WDC also recognized that any public health effort to address the risk factors of cognitive decline and dementia must be grounded in the scientific evidence and informed by the scientific literature. The WDC requested the Alzheimer's Association evaluate and report on the state of the evidence on modifiable risk factors for cognitive decline and dementia to support the WDC in any future recommendations.

The Association's task was not to conduct an independent review of all published literature related to risk reduction, but to evaluate the existing reviews, briefly summarize the findings about the existing body of published evidence, and draw conclusions about the current state of the science. The Alzheimer's Association began by reviewing the detailed reviews prepared by the UK Health Forum for the Blackfriars Consensus [4], Alzheimer's Australia [5,6], Alzheimer's Disease International [7], and Deborah Barnes, PhD, and Kristine Yaffe, MD [8]. Of the articles cited in these reviews, the Association paid particular attention to meta-analyses, systematic reviews, and Cochrane reviews; in addition, the Association evaluated more recently published studies on specific modifiable risk factors. The Association consulted with more than a dozen leading researchers and experts in the dementia risk reduction field—both as part of a pre-existing effort on risk reduction and specifically for this effort—to obtain their input on the current state of the science and the completeness and accuracy of our summary and conclusions.

The Alzheimer's Association—from both a scientific and population-based perspective—weighed the evidence for cognitive decline and all-cause dementia based on the consistency of previous reviews, meta-analyses, and scientific peer-reviewed publications; the number and strength of individual studies (including the number of participants, duration of the study, and diversity of the participants); and the types of those studies (prospective, longitudinal, observational, or randomized controlled trials). The summary of the Association's evaluation was presented at the October 2014 WDC meeting and is presented in this report. Since the WDC meeting, the Association has reviewed additional, more recently published abstracts and studies, which were added to this report. These studies did not change the original underlying conclusions reported to the WDC.

2. Summary of the evidence of individual risk factors

The greatest risk factors for late-onset “sporadic” Alzheimer's disease and other dementias are age [9–11],

family history [12–15], and genetic susceptibility genes, such as the Apolipoprotein E ϵ 4 allele [16,17]. However, none of these risk factors can be modified by medical interventions or by individual behavior. A 2010 National Institutes of Health (NIH) *State of the Science* conference found insufficient evidence, on a clinical level, to support the association of any modifiable risk factors and Alzheimer's disease [18]. The evidence in many cases (particularly with respect to dementia as opposed to cognitive decline) is inconclusive due in large part to the limited data collected to date and the limited number of clinical studies involving specific interventions.

However, despite the limitations of the literature, looking at analyses and studies since the 2010 NIH *State of the Science* conference and viewing the data from a population-based health perspective rather than a clinical perspective, we believe there is sufficient evidence: (a) to support the association between several modifiable risk factors and a reduced risk for cognitive decline; and (b) to suggest that some modifiable risk factors may be associated with reduced risk of dementia. This report discusses these risk factors. Conclusions are summarized in Figures 1 and 2.

3. Cardiovascular risk factors

3.1. Diabetes

Based on several meta-analyses, systematic reviews, and recent studies, more than a dozen prospective, observational, and longitudinal studies have shown lower cognitive performance and an increase in the risk of dementia among individuals with diabetes; on balance, the association between diabetes and dementia appears strong, but not conclusive [19–26]. Further, a recent meta-analysis demonstrated that individuals with mild cognitive impairment (MCI) and diabetes were more likely to progress to dementia than individuals with MCI and no diabetes [27]. Some evidence suggests diabetes increases dementia risk not only through vascular pathways but also through interactions of other biological mechanisms related to diabetes itself [28–30].

3.2. Mid-life obesity

Based on several meta-analyses, systematic reviews, and individual studies, evidence from at least a half dozen prospective studies found that mid-life obesity is associated with an increased risk of dementia. Most postulate this is a strong link, especially with regard to cognitive decline [20,22,31–36]. The association may change with age, as being overweight—and, even possibly being obese—in later life has been associated with reduced risk of dementia [37–41]. And, a recent, large, retrospective cohort study found a lower risk for dementia among those who were overweight even in midlife, while those who were underweight had an elevated risk [42].




Fig. 1. Strength of evidence on risk factors for cognitive decline.

3.3. Mid-life hypertension

Meta-analyses of clinical trials and prospective, observational, longitudinal, and cross-sectional studies, including a Cochrane review, have not indicated a consistent relationship between high blood pressure and dementia; there is stronger evidence for a link with cognitive decline [43–47]. A systematic review of meta-analyses, observational studies, and randomized controlled trials found treatments of hypertension may reduce the risk of cognitive decline [48]; a meta-analysis of longitudinal studies concluded the opposite [49].

Similar to data on the link between obesity and cognitive decline/dementia, studies demonstrate that later-life hypertension may be protective against cognitive decline [50,51].

3.4. Hyperlipidemia (elevated cholesterol)

Systematic reviews of prospective studies have found mixed results for the relationship between both mid-life and late-life high cholesterol levels and dementia, including no association between cholesterol levels and vascular dementia [52,53]. While some observational studies have




Fig. 2. Strength of evidence on risk factors for dementia.

suggested that statin medications used to control cholesterol levels may reduce the risk of dementia [44,54,55], a Cochrane review and systematic reviews found no or inconsistent evidence that use of statins reduces risk [56–58]. The effect has not been seen to date in trials and high-quality cohort studies.

4. Lifestyle risk factors

4.1. Current smoking

According to several systematic reviews and meta-analyses, prospective and longitudinal studies have found strong evidence that current smoking increases the risk of cognitive decline and possibly dementia [7,26,35,59–64]. Quitting smoking may reduce the associated risk to levels comparable to those who have not smoked [7,63–65]. One study of a large multi-ethnic cohort found heavy smoking in middle-age as much as doubled the risk of later-life dementia [66].

4.2. Physical activity

According to systematic reviews and meta-analyses, more than 20 prospective, longitudinal, and cross-sectional studies, as well as randomized controlled trials, have shown physical activity—even in some cases, mild physical activity such as walking—is associated with a decreased risk of cognitive impairment and/or improved cognitive function [35,62,67–76]. Several randomized controlled trials and a Cochrane review of such trials have found that inactive, but otherwise healthy, seniors who begin an exercise program experience significantly improved cognitive function [77,78]. Studies most consistently demonstrate the exercise must be regular and tend toward the more vigorous side [69–72,74]; however, to date, they have failed to pinpoint the optimal duration of the activity, the type and intensity of the exercise, and what period during a person's lifespan it should occur that would maximize potential protective effects.

4.3. Diet

Information on the effects of various aspects of diet (including various nutrients and vitamins, foods, or food groups) on reducing risk is limited and conflicting. Given that many elements of diet are interrelated and interactive, the idea of a whole dietary pattern approach has gained some ground. However, interpretation is challenging as dietary pattern often varies with other lifestyle factors and with demographic variables that may also have an impact on risk. A few cohort studies on the Mediterranean diet (relatively little red meat with an emphasis on whole grains, fruits and vegetables, fish, nuts, and olive oil) or a combined Mediterranean-DASH (Dietary Approaches to Stop Hypertension) diet suggest an association between these diets and reduced risk [79–81].

4.4. Alcohol

Meta-analyses of prospective and case control studies of older adults suggest small or moderate alcohol consumption by older individuals may decrease the risk of cognitive decline and dementia [82–84]. The evidence is not strong enough, however, to suggest those who do not drink should start drinking, especially when weighed against the potential negative effects of excessive alcohol consumption, such as an increased risk of falls among older adults [85–87].

4.5. Cognitive training

A Cochrane review found three dozen randomized controlled trials of mental engagement/cognitive training interventions showing improvements in immediate and delayed recall among those in the treatment group compared with the control group [88]. Systematic reviews of observational studies and randomized controlled trials reached similar conclusions [89,90]. However, based on these analyses, it is unclear whether the improvement is attributed specifically to the cognitive intervention. Despite the large number of trials, most were fairly small, and the data overall were inconclusive. And, as with physical activity, the “recipe” for any successful engagement remains unknown.

4.6. Social engagement

There are very few systematic reviews of the evidence on social engagement—such as doing volunteer work, joining a club, or going to church—as a potential protective factor against cognitive decline or dementia [91]. Some individual studies have shown that social activities, larger social networks, and a history of social contact are associated with better cognitive function and reduced risk for cognitive decline [92–101]. However, an independent coordinated analysis of four longitudinal studies found no effect on cognitive functioning [102]. Looking at the totality of the evidence, most studies in this area are small, are combined with cognitive training and/or physical activities (making it difficult to disaggregate the potential benefits solely of social engagement), and/or are too dissimilar in types of social engagement to draw any conclusions.

5. Other risk factors

5.1. Years of formal education

Among potentially modifiable risk factors, the most consistent evidence surrounds years of formal education (years of schooling in a classroom-based setting taught by professionally trained teachers). People with more years of formal education (measured by grade level attained and/or college attendance) or greater literacy have a lower risk for dementia than those with fewer years of formal education [62,103–111].

5.2. Traumatic brain injury

Solid evidence exists that moderate and severe traumatic brain injury (TBI) increases the risk of developing certain forms of dementia [112–117]. And those who experience repeated head injuries (such as boxers, football players, and combat veterans) may be at an even higher risk [118–124]. While it is not known what specific aspect of TBI (force, repetitiveness, etc.) leads to disrupted brain function, these multiple studies taken together strongly link TBI to increased risk of cognitive decline and dementia.

5.3. Depression

Meta-analyses of cohort and longitudinal studies, as well as additional cohort studies, have shown a history of depression increases the risk for dementia [125–127]. While a recent cohort study found depressive symptoms are independently associated with cognitive decline [128], questions remain regarding whether depression may increase an individual's risk or be an early marker of brain changes associated with dementia. In addition, the effect of treatment for depression on subsequent cognitive functioning is not well understood.

5.4. Sleep

Several cohort and observational studies link sleep disturbances (for example, insomnia and sleep apnea) to increased risk for cognitive decline [129–137]. A recent study further suggested that treatment for breathing disorders that occur during sleep—specifically with continuous positive airway pressure (CPAP)—may reduce the risk of cognitive decline [134]. However, how the exact nature or duration of an individual's sleep problems are related to increased risk is not well understood, nor is it clear whether the sleep disturbances are a cause of or a related precursor to dementia.

6. A multivariate approach to risk reduction

Recent data from several retrospective studies have found declining dementia prevalence or incidence rates in specific population cohorts since the 1970s [138–142]. Some of these studies have noted the large improvements in educational attainment (including higher rates of graduation from high school and college attendance) among the more recent cohorts [138,139,142], underscoring the existing evidence that formal education is beneficial for reducing an individual's risk for cognitive decline and dementia [62,103–111]. Some studies have also noted that over the same period of declining dementia prevalence or incidence, cohorts of study participants had substantial improvements in management of cardiovascular risk factors as well as considerable declines in smoking, heart disease, and stroke [138–140,142]. This has led to the suggestion that medical and public health interventions to lower various cardiovascular risk factors may have had the

additional benefit of improving cognitive health in those specific populations [143,144]. Furthermore, some studies have shown those with multiple vascular issues are at higher risk for cognitive decline and each additional vascular risk factor may be additive to the individual's risk of cognitive decline [145,146].

Given that most of these cardiovascular risk factors are interrelated and connected to the general notion of a healthy lifestyle, focusing on one single lifestyle or health factor may be insufficient to reduce an individual's risk of developing cognitive decline and/or dementia. Some have suggested the most effective strategy may be to address multiple risk factors simultaneously [145,147,148]. The idea of a multivariate approach to reducing risk for cognitive decline was tested in the Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER)—the first published randomized controlled trial on multivariate risk factors and cognitive decline. Recent results from the study, which selected subjects with higher cardiovascular risk profiles, found overall cognitive performance as well as executive function improved significantly with a multi-component lifestyle intervention involving physical activity, nutritional guidance, cognitive training, social activities, and management of cardiovascular risk factors [149].

7. Conclusions

As a result of the Association's review of the current state of evidence on risk factors related to both cognitive decline and dementia, it is clear that there are still many unanswered questions and significant uncertainty with respect to the relationship between individual risk factors and dementia (for example, to what degree there is a causal relationship). There is a clear need for more research on risk reduction, prevention, and brain health—both more longitudinal, population-based cohort studies and randomized controlled trials on the effectiveness of specific interventions that address modifiable risk factors.

However, the Association also believes there is sufficiently strong evidence, from a population-based perspective, to conclude: (1) regular physical activity and management of cardiovascular risk factors (diabetes, obesity, smoking, and hypertension) have been shown to reduce the risk of cognitive decline and may reduce the risk of dementia; and (2) a healthy diet and lifelong learning/cognitive training may also reduce the risk of cognitive decline. The Institute of Medicine, drawing on a panel of distinguished researchers in the field, recently examined the evidence with regard to cognitive aging and cognitive decline and reached a virtually identical conclusion [150].

The evidence has now reached a point that it can no longer remain simply an exercise in academic discussion. The public should know what the science concludes: certain healthy behaviors known to be effective for diabetes,

cardiovascular disease, and cancer are also good for brain health and for reducing the risk of cognitive decline. For our part, the Alzheimer's Association is launching a new brain health education program, *Healthy Habits for a Healthier You*. It is designed to provide consumers with the latest research and practical information on ways they can take care of their bodies and brains to age as well as possible.

For maximum impact, a broader effort must be undertaken—and governments must be involved—to increase public awareness and education about known and potentially modifiable risk factors of cognitive decline and dementia. Since the Association presented its conclusions, the WDC issued a statement encouraging all nations not only to invest in further research, but also to incorporate dementia risk reduction and management into public health policies, public health campaigns, and non-communicable disease strategies and action plans [151]. It is a two-pronged approach: promote risk reduction today based on the scientific evidence available and pursue more research so the evidence will be stronger and more definitive tomorrow.

References

- [1] Prince M, Guerchet M, Prina M. *The Epidemiology and Impact of Dementia: Current State and Future Trends*. Geneva: World Health Organization; 2015.
- [2] Prince M, Guerchet M, Prina M, Alzheimer's Disease International. *Policy Brief for Heads of Government: The Global Impact of Dementia 2013-2050*. London: Alzheimer's Disease International; 2013.
- [3] G8 Dementia Summit Communique. Available at: www.gov.uk/government/uploads/system/uploads/attachment_data/file/265868/2901669_G8_DementiaSummitCommunique_acc.pdf. Accessed on April 7, 2015.
- [4] UK Health Forum. *Promoting brain health: Developing a prevention agenda linking dementia and other non-communicable diseases*. London: UK Health Forum; 2014.
- [5] Woodward M. *Dementia Risk Reduction: The Evidence*. Scullin, Australia: Alzheimer's Australia; 2007. Paper 13.
- [6] Farrow M, O'Connor E. *Targeting Brain, Body and Heart for Cognitive Health and Dementia Prevention: Current Evidence and Future Directions*. Canberra, Australia: Alzheimer's Australia; 2012. Paper 29.
- [7] Prince M, Albanese E, Guerchet M, Prina M. *Dementia and Risk Reduction: An Analysis of Protective and Modifiable Factors*. London: Alzheimer's Disease International; 2014. World Alzheimer's Report 2014.
- [8] Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol* 2011;10:819–28.
- [9] Alzheimer's Association. *2014 Alzheimer's Disease Facts and Figures*. *Alzheimers Dement* 2014;10:e47–92.
- [10] Hebert LE, Bienias JL, Aggarwal NT, Wilson RS, Bennett DA, Shah RC, et al. Change in risk of Alzheimer disease over time. *Neurology* 2010;75:786–91.
- [11] Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology* 2013;80:1778–83.
- [12] Green RC, Cupples LA, Go R, Benke KS, Edeki T, Griffith PA, et al. Risk of dementia among white and African American relatives of patients with Alzheimer disease. *JAMA* 2002;287:329–36.
- [13] Fratiglioni L, Ahlborn A, Viitanen M, Winblad B. Risk factors for late-onset Alzheimer's disease: A population-based, case-control study. *Ann Neurol* 1993;33:258–66.
- [14] Mayeux R, Sano M, Chen J, Tatemichi T, Stern Y. Risk of dementia in first-degree relatives of patients with Alzheimer's disease and related disorders. *Arch Neurol* 1991;48:269–73.
- [15] Lautenschlager NT, Cupples LA, Rao VS, Auerbach SA, Becker R, Burke J, et al. Risk of dementia among relatives of Alzheimer's disease patients in the MIRAGE Study: What is in store for the oldest old? *Neurology* 1996;46:641–50.
- [16] Saunders AM, Strittmatter WJ, Schmechel D, George-Hyslop PH, Pericak-Vance MA, Joo SH, et al. Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology* 1993;43:1467–72.
- [17] Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease: A meta-analysis. *JAMA* 1997;278:1349–56.
- [18] National Institutes of Health. *NIH Consensus Development Conference Statement on Preventing Alzheimer's Disease and Cognitive Decline*. Bethesda, MD: National Institutes of Health; 2010.
- [19] Lu FP, Lin KP, Kuo HK. Diabetes and the risk of multi-system aging phenotypes: A systematic review and meta-analysis. *PLoS One* 2009;4:e4144.
- [20] Profenno LA, Porsteinsson AP, Faraone SV. Meta-analysis of Alzheimer's disease risk with obesity, diabetes, and related disorders. *Biol Psychiatry* 2010;67:505–12.
- [21] Biessels GJ, Staekenborg S, Brunner E, Brayne C, Scheltens P. Risk of dementia in diabetes mellitus: A systematic review. *Lancet Neurol* 2006;5:64–74.
- [22] Kloppenborg RP, van den Berg E, Kappelle LJ, Biessels GJ. Diabetes and other vascular risk factors for dementia: Which factor matters most? A systematic review. *Eur J Pharmacol* 2008;585:97–108.
- [23] Cheng G, Huang C, Deng H, Wang H. Diabetes as a risk factor for dementia and mild cognitive impairment: A meta-analysis of longitudinal studies. *Intern Med J* 2012;42:484–91.
- [24] Gudala K, Bansal D, Schifano F, Bhansali A. Diabetes mellitus and risk of dementia: A meta-analysis of prospective observational studies. *J Diabetes Investig* 2013;4:640–50.
- [25] Roberts RO, Knopman DS, Cha RH, Mielke MM, Pankratz VS, Boeve BF, et al. Diabetes and elevated hemoglobin A1c levels are associated with brain hypometabolism but not amyloid accumulation. *J Nucl Med* 2014;55:759–64.
- [26] Plassman BL, Williams JW Jr, Burke JR, Holsinger T, Benjamin S. Systematic review: factors associated with risk for and possible prevention of cognitive decline in later life. *Ann Intern Med* 2010;153:182–93.
- [27] Cooper C, Sommerlad A, Lyketsos CG, Livingston G. Modifiable predictors of dementia in mild cognitive impairment: a systematic review and meta-analysis. *Am J Psychiatry* 2015;172:323–34.
- [28] Mushtaq G, Khan JA, Kamal MA. Biological mechanisms linking Alzheimer's disease and Type-2 Diabetes mellitus. *CNS Neurol Disord Drug Targets* 2014;13:1192–201.
- [29] De Felice FG, Ferreira ST. Inflammation, defective insulin signaling, and mitochondrial dysfunction as common molecular denominators connecting type 2 diabetes to Alzheimer disease. *Diabetes* 2014;63:2262–72.
- [30] Yang Y, Song W. Molecular links between Alzheimer's disease and diabetes mellitus. *Neuroscience* 2013;250:140–50.
- [31] Sellbom KS, Gunstad J. Cognitive function and decline in obesity. *J Alzheimers Dis* 2012;30(Suppl 2):S89–95.
- [32] Beydoun MA, Beydoun HA, Wang Y. Obesity and central obesity as risk factors for incident dementia and its subtypes: A systematic review and meta-analysis. *Obes Rev* 2008;9:204–18.
- [33] Loeff M, Walach H. Midlife obesity and dementia: Meta-analysis and adjusted forecast of dementia prevalence in the United States and China. *Obesity (Silver Spring)* 2013;21:E51–5.

- [34] Anstey KJ, Cherbuin N, Budge M, Young J. Body mass index in midlife and late-life as a risk factor for dementia: A meta-analysis of prospective studies. *Obes Rev* 2011;12:e426–37.
- [35] Lee Y, Back JH, Kim J, Kim SH, Na DL, Cheong HK, et al. Systematic review of health behavioral risks and cognitive health in older adults. *Int Psychogeriatr* 2010;22:174–87.
- [36] Gustafson DR, Backman K, Joas E, Waern M, Ostling S, Guo X, et al. 37 years of body mass index and dementia: Observations from the prospective population study of women in Gothenburg, Sweden. *J Alzheimers Dis* 2012;28:163–71.
- [37] Luchsinger JA, Patel B, Tang MX, Schupf N, Mayeux R. Measures of adiposity and dementia risk in elderly persons. *Arch Neurol* 2007;64:392–8.
- [38] Fitzpatrick AL, Kuller LH, Lopez OL, Diehr P, O'Meara ES, Longstreth WT Jr, et al. Midlife and late-life obesity and the risk of dementia: Cardiovascular health study. *Arch Neurol* 2009;66:336–42.
- [39] Barnes DE, Covinsky KE, Whitmar RA, Kuller LH, Lopez OL, Yaffe K. Predicting risk of dementia in older adults: the late-life dementia risk index. *Neurology* 2009;73:173–9.
- [40] Dahl AK, Lopponen M, Isoaho R, Berg S, Kivela SL. Overweight and obesity in old age are not associated with greater dementia risk. *J Am Geriatr Soc* 2008;56:2261–6.
- [41] Gustafson DR, Luchsinger JA. High adiposity: Risk factor for dementia and Alzheimer's disease? *Alzheimers Res Ther* 2013;5:57.
- [42] Qizilbash N, Gregson J, Johnson ME, Pearce N, Douglas I, Wing K, et al. BMI and risk of dementia in two million people over two decades: a retrospective cohort study. *Lancet Diabetes Endocrinol* 2015; [http://dx.doi.org/10.1016/S2213-8587\(15\)00033-9](http://dx.doi.org/10.1016/S2213-8587(15)00033-9). Epub ahead of print.
- [43] Power MC, Weuve J, Gagne JJ, McQueen MB, Viswanathan A, Blacker D. The association between blood pressure and incident Alzheimer disease: A systematic review and meta-analysis. *Epidemiology* 2011;22:646–59.
- [44] McGuinness B, Todd S, Passmore P, Bullock R. Blood pressure lowering in patients without prior cerebrovascular disease for prevention of cognitive impairment and dementia. *Cochrane Database Syst Rev* 2009;(4):CD004034.
- [45] Shah K, Qureshi SU, Johnson M, Parikh N, Schulz PE, Kunik ME. Does use of antihypertensive drugs affect the incidence or progression of dementia? A systematic review. *Am J Geriatr Pharmacother* 2009;7:250–61.
- [46] Guan JW, Huang CQ, Li YH, Wan CM, You C, Wang ZR, et al. No association between hypertension and risk for Alzheimer's disease: A meta-analysis of longitudinal studies. *J Alzheimers Dis* 2011;27:799–807.
- [47] Sharp SI, Aarsland D, Day S, Sonnesyn H, Ballard C, Alzheimer's Society Vascular Dementia Systematic Review Group. Hypertension is a potential risk factor for vascular dementia: Systematic review. *Int J Geriatr Psychiatry* 2011;26:661–9.
- [48] Rouch L, Cestac P, Hanon O, Cool C, Helmer C, Bouhanick B, et al. Antihypertensive drugs, prevention of cognitive decline and dementia: a systematic review of observational studies, randomized controlled trials and meta-analyses, with discussion of potential mechanisms. *CNS Drugs* 2015;29:113–30.
- [49] Chang-Quan H, Hui W, Chao-Min W, Zheng-Rong W, Jun-Wen G, Yong-Hong L, et al. The association of antihypertensive medication use with risk of cognitive decline and dementia: a meta-analysis of longitudinal studies. *Int J Clin Pract* 2011;65:1295–305.
- [50] Kennelly SP, Lawlor BA, Kenny RA. Blood pressure and the risk for dementia: A double edged sword. *Ageing Res Rev* 2009;8:61–70.
- [51] Corrada M, Hayden KM, Bullain SS, Paganini-Hill A, DeMoss J, Aguirre C, et al. Age of onset of hypertension and risk of dementia in the oldest-old: The 90+ study. *Alzheimers Dement* 2014;10(4 Supplement):P501.
- [52] Anstey KJ, Lipnicki DM, Low LF. Cholesterol as a risk factor for dementia and cognitive decline: A systematic review of prospective studies with meta-analysis. *Am J Geriatr Psychiatry* 2008;16:343–54.
- [53] Kivipelto M, Solomon A. Cholesterol as a risk factor for Alzheimer's disease – epidemiological evidence. *Acta Neurol Scand Suppl* 2006;185:50–7.
- [54] Muangpaisan W, Brayne C, Alzheimer's Society Vascular Dementia Systematic Review Group. Systematic review of statins for the prevention of vascular dementia or dementia. *Geriatr Gerontol Int* 2010;10:199–208.
- [55] Beri A, Sural N, Mahajan SB. Non-atheroprotective effects of statins: A systematic review. *Am J Cardiovasc Drugs* 2009;9:361–70.
- [56] McGuinness B, Craig D, Bullock R, Passmore P. Statins for the prevention of dementia. *Cochrane Database Syst Rev* 2009;(2):CD003160.
- [57] Richardson K, Schoen M, French B, Umscheid CA, Mitchell MD, Arnold SE, et al. Statins and cognitive function: a systematic review. *Ann Intern Med* 2013;159:688–97.
- [58] Ligthart SA, Moll van Charante EP, Van Gool WA, Richard E. Treatment of cardiovascular risk factors to prevent cognitive decline and dementia: a systematic review. *Vasc Health Risk Manag* 2010;6:775–85.
- [59] Anstey KJ, von Sanden C, Salim A, O'Kearney R. Smoking as a risk factor for dementia and cognitive decline: A meta-analysis of prospective studies. *Am J Epidemiol* 2007;166:367–78.
- [60] Cataldo JK, Prochaska JJ, Glantz SA. Cigarette smoking is a risk factor for Alzheimer's disease: An analysis controlling for tobacco industry affiliation. *J Alzheimers Dis* 2010;19:465–80.
- [61] Peters R, Poulter R, Warner J, Beckett N, Burch L, Bulpitt C. Smoking, dementia and cognitive decline in the elderly: A systematic review. *BMC Geriatr* 2008;8:36.
- [62] Beydoun MA, Beydoun HA, Gamaldo AA, Teel A, Zonderman AB, Wang Y. Epidemiologic studies of modifiable factors associated with cognition and dementia: Systematic review and meta-analysis. *BMC Public Health* 2014;14:643.
- [63] McKenzie J, Bhatti L, Tursan d'Espaignet E. WHO Tobacco Knowledge Summaries: Tobacco and Dementia. Geneva: World Health Organization; 2014.
- [64] Zhong G, Wang Y, Zhang Y, Guo JJ, Zhao Y. Smoking is associated with an increased risk of dementia: A meta-analysis of prospective cohort studies with investigation of potential effect modifiers. *PLoS One* 2015;10:e0118333.
- [65] Sabia S, Elbaz A, Dugravot A, Head J, Shipley M, Hagger-Johnson G, et al. Impact of smoking on cognitive decline in early old age: The Whitehall II cohort study. *Arch Gen Psychiatry* 2012;69:627–35.
- [66] Rusanen M, Kivipelto M, Quesenberry CP Jr, Zhou J, Whitmer RA. Heavy smoking in midlife and long-term risk of Alzheimer disease and vascular dementia. *Arch Intern Med* 2011;171:333–9.
- [67] Rolland Y, Abellan van Kan G, Vellas B. Physical activity and Alzheimer's disease: From prevention to therapeutic perspectives. *J Am Med Dir Assoc* 2008;9:390–405.
- [68] Hamer M, Chida Y. Physical activity and risk of neurodegenerative disease: A systematic review of prospective evidence. *Psychol Med* 2009;39:3–11.
- [69] Paterson DH, Warburton DE. Physical activity and functional limitations in older adults: A systematic review related to Canada's Physical Activity Guidelines. *Int J Behav Nutr Phys Act* 2010;7:38.
- [70] Lautenschlager NT, Cox KL, Flicker L, Foster JK, van Bockxmeer FM, Xiao J, et al. Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: A randomized trial. *JAMA* 2008;300:1027–37.
- [71] Sofi F, Valecchi D, Bacci D, Abbate R, Gensini GF, Casini A, et al. Physical activity and risk of cognitive decline: A meta-analysis of prospective studies. *J Intern Med* 2011;269:107–17.
- [72] Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia? A systematic review

- and meta-analysis of longitudinal studies. *BMC Public Health* 2014; 14:510.
- [73] Ahlskog JE, Geda YE, Graff-Radford NR, Petersen RC. Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. *Mayo Clin Proc* 2011;86:876–84.
- [74] Colcombe S, Kramer AF. Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci* 2003;14:125–30.
- [75] Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K, et al. Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosom Med* 2010;72:239–52.
- [76] Bherer L, Erickson KI, Liu-Ambrose T. A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J Aging Res* 2013;2013:657508.
- [77] Angevaren M, Aufdemkampe G, Verhaar HJ, Aleman A, Vanhees L. Physical activity and enhanced fitness to improve cognitive function in older people without known cognitive impairment. *Cochrane Database Syst Rev* 2008;(3):CD005381.
- [78] Barnes DE, Santos-Modesitt W, Poelke G, Kramer AF, Castro C, Middleton LE, et al. The Mental Activity and eXercise (MAX) trial: A randomized controlled trial to enhance cognitive function in older adults. *JAMA Intern Med* 2013;173:797–804.
- [79] Lourida I, Soni M, Thompson-Coon J, Purandare N, Lang IA, Ukoumunne OC, et al. Mediterranean diet, cognitive function, and dementia: A systematic review. *Epidemiology* 2013;24:479–89.
- [80] Psaltopoulou T, Sergentanis TN, Panagiotakos DB, Sergentanis IN, Kosti R, Scarmeas N. Mediterranean diet, stroke, cognitive impairment, and depression: A meta-analysis. *Ann Neurol* 2013; 74:580–91.
- [81] Morris MC, Tangney CC, Wang Y, Sacks FM, Bennett DA, Aggarwal NT. MIND diet associated with reduced incidence of Alzheimer's disease. *Alzheimers Dement* 2015; <http://dx.doi.org/10.1016/j.jalz.2014.11.009>. Epub ahead of print.
- [82] Anstey KJ, Mack HA, Cherbuin N. Alcohol consumption as a risk factor for dementia and cognitive decline: Meta-analysis of prospective studies. *Am J Geriatr Psychiatry* 2009;17:542–55.
- [83] Peters R, Peters J, Warner J, Beckett N, Bulpitt C. Alcohol, dementia and cognitive decline in the elderly: A systematic review. *Age Ageing* 2008;37:505–12.
- [84] Neafsey EJ, Collins MA. Moderate alcohol consumption and cognitive risk. *Neuropsychiatr Dis Treat* 2011;7:465–84.
- [85] Mukamal KJ, Mittleman MA, Longstreth WT Jr, Newman AB, Fried LP, Siscovick DS. Self-reported alcohol consumption and falls in older adults: Cross-sectional and longitudinal analyses of the Cardiovascular Health Study. *J Am Geriatr Soc* 2004;52:1174–9.
- [86] Stahre M, Roeber J, Kanny D, Brewer RD, Zhang X. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis* 2014;11:E109.
- [87] U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary Guidelines for Americans*, 2010. 7th ed. Washington, DC: U.S. Government Printing Office; 2010.
- [88] Martin M, Clare L, Altgassen AM, Cameron MH, Zehnder F. Cognition-based interventions for healthy older people and people with mild cognitive impairment. *Cochrane Database Syst Rev* 2011;(1):CD006220.
- [89] Stern C, Munn Z. Cognitive leisure activities and their role in preventing dementia: A systematic review. *Int J Evid Based Healthc* 2010;8:2–17.
- [90] Valenzuela M, Sachdev P. Can cognitive exercise prevent the onset of dementia? Systematic review of randomized clinical trials with longitudinal follow-up. *Am J Geriatr Psychiatry* 2009;17:179–87.
- [91] Fratiglioni L, Paillard-Borg S, Winblad B. An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol* 2004;3:343–53.
- [92] James BD, Wilson RS, Barnes LL, Bennett DA. Late-life social activity and cognitive decline in old age. *J Int Neuropsychol Soc* 2011; 17:998–1005.
- [93] Noice T, Noice H, Kramer AF. Participatory arts for older adults: a review of benefits and challenges. *Gerontologist* 2014;54:741–53.
- [94] Ertel KA, Glymour MM, Berkman LF. Effects of social integration on preserving memory function in a nationally representative US elderly population. *Am J Public Health* 2008;98:1215–20.
- [95] Stine-Morrow EA, Parisi JM, Morrow DG, Park DC. The effects of an engaged lifestyle on cognitive vitality: A field experiment. *Psychol Aging* 2008;23:778–86.
- [96] Karp A, Paillard-Borg S, Wang HX, Silverstein M, Winblad B, Fratiglioni L. Mental, physical and social components in leisure activities equally contribute to decrease dementia risk. *Dement Geriatr Cogn Disord* 2006;21:65–73.
- [97] Barnes LL, Mendes de Leon CF, Wilson RS, Bienias JL, Evans DA. Social resources and cognitive decline in a population of older African Americans and whites. *Neurology* 2004;63:2322–6.
- [98] Holtzman RE, Rebok GW, Saczynski JS, Kouzis AC, Wilcox Doyle K, Eaton WW. Social network characteristics and cognition in middle-aged and older adults. *J Gerontol B Psychol Sci Soc Sci* 2004;59:P278–84.
- [99] Scarmeas N, Levy G, Tang MX, Manly J, Stern Y. Influence of leisure activity on the incidence of Alzheimer's disease. *Neurology* 2001; 57:2236–42.
- [100] Crooks VC, Lubben J, Petitti DB, Little D, Chiu V. Social network, cognitive function, and dementia incidence among elderly women. *Am J Public Health* 2008;98:1221–7.
- [101] Seeman TE, Miller-Martinez DM, Stein Merkin S, Lachman ME, Tun PA, Karlamangla AS. Histories of social engagement and adult cognition: Midlife in the U.S. study. *J Gerontol B Psychol Sci Soc Sci* 2011;66(Supplement 1):i141–52.
- [102] Brown CL, Gibbons LE, Kennison RF, Robitaille A, Lindwall M, Mitchell MB, et al. Social activity and cognitive functioning over time: a coordinated analysis of four longitudinal studies. *J Aging Res* 2012;2012:287438.
- [103] Meng X, D'Arcy C. Education and dementia in the context of the cognitive reserve hypothesis: a systematic review with meta-analyses and qualitative analyses. *PLoS One* 2012;7:e38268.
- [104] Jefferson AL, Gibbons LE, Rentz DM, Carvalho JO, Manly J, Bennett DA, et al. A life course model of cognitive activities, socioeconomic status, education, reading ability, and cognition. *J Am Geriatr Soc* 2011;59:1403–11.
- [105] Fitzpatrick AL, Kuller LH, Ives DG, Lopez OL, Jagust W, Breitner JC, et al. Incidence and prevalence of dementia in the Cardiovascular Health Study. *J Am Geriatr Soc* 2004;52:195–204.
- [106] Kukull WA, Higdon R, Bowen JD, McCormick WC, Teri L, Schellenberg GD, et al. Dementia and Alzheimer disease incidence: A prospective cohort study. *Arch Neurol* 2002;59:1737–46.
- [107] Evans DA, Bennett DA, Wilson RS, Bienias JL, Morris MC, Scherr PA, et al. Incidence of Alzheimer disease in a biracial urban community: Relation to apolipoprotein E allele status. *Arch Neurol* 2003;60:185–9.
- [108] Stern Y, Gurland B, Tatemichi TK, Tang MX, Wilder D, Mayeux R. Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA* 1994;271:1004–10.
- [109] Evans DA, Hebert LE, Beckett LA, Scherr PA, Albert MS, Chown MJ, et al. Education and other measures of socioeconomic status and risk of incident Alzheimer disease in a defined population of older persons. *Arch Neurol* 1997;54:1399–405.
- [110] Sando SB, Melquist S, Cannon A, Hutton M, Sletvoid O, Saltvedt I, et al. Risk-reducing effect of education in Alzheimer's disease. *Int J Geriatr Psychiatry* 2008;23:1156–62.
- [111] Caamano-Isorna F, Corral M, Montes-Martinez A, Takkouche B. Education and dementia: a meta-analytic study. *Neuroepidemiology* 2006;26:226–32.
- [112] Lye TC, Shores EA. Traumatic brain injury as a risk factor for Alzheimer's disease: A review. *Neuropsychol Rev* 2000;10:115–29.
- [113] Plassman BL, Havlik RJ, Steffens DC, Helms MJ, Newman TN, Drosdick D, et al. Documented head injury in early adulthood and

- risk of Alzheimer's disease and other dementias. *Neurology* 2000; 55:1158–66.
- [114] Barnes DE, Kaup A, Kirby KA, Byers AL, Diaz-Arrastia R, Yaffe K. Traumatic brain injury and risk of dementia in older veterans. *Neurology* 2014;83:312–9.
- [115] Gardner R, Yaffe K. Traumatic brain injury may increase risk of young onset dementia. *Ann Neurol* 2014;75:339–41.
- [116] Fleminger S, Oliver DL, Lovestone S, Rabe-Hesketh S, Giora A. Head injury as a risk factor for Alzheimer's disease: the evidence 10 years on; a partial replication. *J Neurol Neurosurg Psychiatry* 2003;74:857–62.
- [117] Shively S, Scher AI, Perl DP, Diaz-Arrastia R. Dementia resulting from traumatic brain injury: what is the pathology? *Arch Neurol* 2012;69:1245–51.
- [118] Smith DH, Johnson VE, Stewart W. Chronic neuropathologies of single and repetitive TBI: Substrates of dementia? *Nat Rev Neuro* 2013; 9:211–21.
- [119] Guskiewicz KM. Association between recurrent concussion and late-life cognitive impairment in retired professional football players. *Neurosurgery* 2005;57:719–26.
- [120] Institute for Social Research. National Football League Player Care Foundation Study of NFL Retired Players. Ann Arbor: University of Michigan; 2009.
- [121] Lehman EJ, Hein MJ, Baron SL, Gersix CM. Neurodegenerative causes of death among retired National Football League players. *Neurology* 2012;79:1970–4.
- [122] Groswasser Z, Reider-Groswasser II, Schwab K, Ommaya AK, Pridgen A, Brown HR, et al. Quantitative imaging in late TBI. Part II: Cognition and work after closed and penetrating head injury: A report of the Vietnam Head Injury Study. *Brain Inj* 2002;16:681–90.
- [123] Salazar AM, Warden DL, Schwab K, Spector J, Braverman S, Walter J, et al. Cognitive rehabilitation for traumatic brain injury: A randomized trial. Defense and Veterans Head Injury Program (DVHIP) Study Group. *JAMA* 2000;283:3075–81.
- [124] Roberts GW, Allsop D, Bruton C. The occult aftermath of boxing. *J Neurol Neurosurg Psychiatry* 1990;53:373–8.
- [125] Ownby RL, Crocco E, Acevedo A, John V, Loewenstein D. Depression and risk for Alzheimer disease: Systematic review, meta-analysis, and meta-regression analysis. *Arch Gen Psychiatry* 2006;63:530–8.
- [126] Diniz BS, Butters MA, Albert SM, Dew MA, Reynolds CF 3rd. Late-life depression and risk of vascular dementia and Alzheimer's disease: Systematic review and meta-analysis of community-based cohort studies. *Br J Psychiatry* 2013;202:329–35.
- [127] Barnes DE, Yaffe K, Byers AL, McCormick M, Schaefer C, Whitmer RA. Midlife vs late-life depressive symptoms and risk of dementia: Differential effects for Alzheimer disease and vascular dementia. *Arch Gen Psychiatry* 2012;69:493–8.
- [128] Wilson RS, Capuano AW, Boyle PA, Hoganson GM, Hibel LP, Shah RC, et al. Clinical-pathologic study of depressive symptoms and cognitive decline in old age. *Neurology* 2014;83:702–9.
- [129] Yaffe K, Laffan AM, Harrison SL, Redline S, Spira AP, Ensrud KE, et al. Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA* 2011;306:613–9.
- [130] Ju YE, Lucey BP, Holtzman DM. Sleep and Alzheimer disease pathology—a bidirectional relationship. *Nat Rev Neuro* 2014;10:115–9.
- [131] Chang WP, Liu ME, Chang WC, Yang AC, Ku YC, Pai JT, et al. Sleep apnea and the risk of dementia: A population-based 5-year follow-up study in Taiwan. *PLoS One* 2013;8:e78655.
- [132] Chen PL, Lee WJ, Sun WZ, Oyang YJ, Fuh JL. Risk of dementia in patients with insomnia and long-term use of hypnotics: A population-based retrospective cohort study. *PLoS One* 2012;7:e49113.
- [133] Sterniczuk R, Theou O, Rusak B, Rockwood K. Sleep disturbance is associated with incident dementia and mortality. *Curr Alzheimer Res* 2013;10:767–75.
- [134] Osorio RS, Gumb T, Pirraglia E, Varga AW, Lu SE, Lim J, et al. Sleep-disordered breathing advances cognitive decline in the elderly. *Neurology* 2015;84:1964–71.
- [135] Yaffe K, Falvey CM, Hoang T. Connections between sleep and cognition in older adults. *Lancet Neurol* 2014;13:1017–28.
- [136] Lim AS, Kowgier M, Yu L, Buchman AS, Bennett DA. Sleep fragmentation and the risk of incident Alzheimer's disease and cognitive decline in older persons. *Sleep* 2013;36:1027–32.
- [137] Potvin O, Lorrain D, Forget H, Dube M, Grenier S, Preville M, et al. Sleep quality and 1-year incident cognitive impairment in community-dwelling older adults. *Sleep* 2012;35:491–9.
- [138] Satizabal CL, Beiser A, Chene G, Chouraki VA, Himali JJ, Preis SR, et al. Temporal trends in dementia incidence in the Framingham Study. *Alzheimers Dement* 2014;10(4 Supplement):P296.
- [139] Schrijvers EM, Verhaaren BF, Koudstaal PJ, Hofman A, Ikram MA, Breteler MM. Is dementia incidence declining?: Trends in dementia incidence since 1990 in the Rotterdam Study. *Neurology* 2012; 78:1456–63.
- [140] Qiu C, von Strauss E, Backman L, Winblad B, Fratiglioni L. Twenty-year changes in dementia occurrence suggest decreasing incidence in central Stockholm, Sweden. *Neurology* 2013;80:1888–94.
- [141] Dodge HH, Zhu J, Lee CW, Chang CC, Ganguli M. Cohort effects in age-associated cognitive trajectories. *J Gerontol A Biol Sci Med Sci* 2014;69:687–94.
- [142] Doblhammer G, Fink A, Fritze T. Short-term trends in dementia prevalence in Germany between the years 2007 and 2009. *Alzheimers Dement* 2015;11:291–9.
- [143] Elwood P, Galante J, Pickering J, Palmer S, Bayer A, Ben-Shlomo Y, et al. Healthy lifestyles reduce the incidence of chronic diseases and dementia: evidence from the Caerphilly cohort study. *PLoS One* 2013;8:e81877.
- [144] Larson EB, Yaffe K, Langa KM. New insights into the dementia epidemic. *N Engl J Med* 2013;369:2275–7.
- [145] Middleton LE, Yaffe K. Targets for the prevention of dementia. *J Alzheimers Dis* 2010;20:915–24.
- [146] Roberts RO, Cha RH, Mielke MM, Geda YE, Boeve BF, Machulda MM, et al. Risk and protective factors for cognitive impairment in persons aged 85 years and older. *Neurology* 2015;84:1854–61.
- [147] Mangialasche F, Kivipelto M, Solomon A, Fratiglioni L. Dementia prevention: current epidemiological evidence and future perspective. *Alzheimers Res Ther* 2012;4:6.
- [148] Monsuez JJ, Gesquiere-Dando A, Rivera S. Cardiovascular prevention of cognitive decline. *Cardiol Res Pract* 2011;2011:250970.
- [149] Ngandu T, Lehtisalo J, Solomon A, Levalahti E, Ahtiluoto RA, Backman L, et al. A 2-year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. *Lancet* 2015; [http://dx.doi.org/10.1016/S0140-6736\(15\)60461-5](http://dx.doi.org/10.1016/S0140-6736(15)60461-5). Epub ahead of print.
- [150] Institute of Medicine. Cognitive Aging: Progress in Understanding and Opportunity for Action. Washington, DC: The National Academies Press; 2015.
- [151] World Dementia Council. WDC Dementia Risk Reduction Statement. Available at: <https://s3-eu-west-1.amazonaws.com/media.dh.gov.uk/network/353/files/2015/01/WDC-supportive-statement-on-risk-reduction-FINAL-FOR-PUBLICATION-BRANDED-150128-LM.pdf>. Accessed on April 7, 2015.